

HYPOTHESIS TESTING • MTH107

Class R FAQ

by Derek H. Ogle, revised Dec-16

1-Sample Z-Test

$H_0: \mu = \mu_0$ (where $\mu_0 =$ specific value)

Statistic: \bar{X} **Test Statistic:** $Z = \frac{\bar{x} - \mu_0}{\frac{\sigma}{\sqrt{n}}}$ **Conf. Region:** $\bar{X} + Z^* \frac{\sigma}{\sqrt{n}}$

Assumptions: 1) σ is known
2) $n \geq 30$, $n \geq 15$ and popn not strongly skewed, OR popn is normal

R: `z.test()`

1-Sample t-Test

$H_0: \mu = \mu_0$ (where $\mu_0 =$ specific value)

Statistic: \bar{X} **Test Statistic:** $t = \frac{\bar{x} - \mu_0}{\frac{s}{\sqrt{n}}}$ **Conf. Region:** $\bar{X} + t^* \frac{s}{\sqrt{n}}$ **df:** $n-1$

Assumptions: 1) σ is UNknown,
2) $n \geq 40$, $n \geq 15$ & histogram not strongly skewed, OR histogram is normal

R: `t.test()`, `hist()`

2-Sample t-Test

$H_0: \mu_1 = \mu_2$ **Statistic:** $\bar{X}_1 - \bar{X}_2$

Test Statistic: $t = \frac{(\bar{x}_1 - \bar{x}_2) - 0}{\sqrt{s_p^2 \left(\frac{1}{n_1} + \frac{1}{n_2} \right)}}$ where $s_p^2 = \frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2 - 2}$

Conf. Region: $(\bar{x}_1 - \bar{x}_2) + t^* \sqrt{s_p^2 \left(\frac{1}{n_1} + \frac{1}{n_2} \right)}$ **df:** $n_1 + n_2 - 2$

Assumptions:
1) Individuals in popns are independent
2) Variances are equal (use Levene's Test)
3) $n_1 + n_2 \geq 40$, $n_1 + n_2 \geq 15$ and both histograms are not strongly skewed, OR Both histograms are normal

R: `t.test()`, `levenesTest()`, `hist()`

Choosing a Hypothesis Test

1. If response variable is QUANTITATIVE, then GOTO 2; otherwise GOTO 5.

Quantitative Response

2. If 1 POPULATION was sampled, then GOTO 3; otherwise GOTO 4.

3. If σ is KNOWN, then **1-Sample Z**; otherwise **1-Sample t**.

4. If individuals in populations are INDEPENDENT, then **2-Sample t**; otherwise, **Paired t**.

Categorical Response

5. If 1 POPULATION was sampled, then **Goodness-of-Fit**; otherwise, **Chi-Square**.

Making a Decision about H_0

If the p-value $< \alpha$, then **REJECT H_0** , otherwise **DNR H_0** .

Chi-Square Test

H_0 : "Distribution of indivs into response levels is the same for all populations"

H_A : "Distribution of indivs into response levels is NOT the same for all populations"

Statistic: Observed frequency table

Test Statistic: $\chi^2 = \sum \frac{(\text{Observed} - \text{Expected})^2}{\text{Expected}}$ **df:** (rows-1)(columns-1)

Assumptions: ≥ 5 in each cell of the expected table

R: `xtabs()`, `matrix()`, `chisq.test()`, `percTable()`

Goodness-of-Fit Test

H_0 : "Distribution of indivs into response levels follows the theoretical distribution"

H_A : "Distribution of indivs into response levels does NOT follow the theoretical distribution"

Statistic: Observed frequency table

Test Statistic: $\chi^2 = \sum \frac{(\text{Observed} - \text{Expected})^2}{\text{Expected}}$ **df:** cells-1

Assumptions: ≥ 5 in each cell of the expected table

R: `xtabs()`, `c()`, `chisq.test()`, `percTable()`, `chiGOF()`